

Self-Sustained Ionization and Vanishing Dead Zones in Protoplanetary Disks

Shu-ichiro Inutsuka¹ and Takayoshi Sano²

ABSTRACT

We analyze the ionization state of the magnetohydrodynamically turbulent protoplanetary disks and propose a new mechanism of sustaining ionization. First, we show that in the quasi-steady state of turbulence driven by magnetorotational instability in a typical protoplanetary disk with dust grains, the amount of energy dissipation should be sufficient for providing the ionization energy that is required for activating magnetorotational instability. Second, we show that in the disk with dust grains the energetic electrons that compose electric currents in weakly ionized gas can provide collisional ionization, depending on the actual saturation state of magnetorotational turbulence. On the other hand, we show that in the protoplanetary disks with the reduced effect of dust grains, the turbulent motion can homogenize the ionization degree, leading to the activation of magnetorotational instability even in the absence of other ionization processes. The results in this Letter indicate that most of the regions in protoplanetary disks remain magnetically active, and we thus require a change in the theoretical modeling of planet formation.

Subject headings: protoplanetary disks — planetary systems: formation — instabilities — MHD — turbulence — accretion, accretion disks

1. Introduction

Protoplanetary disks are supposed to be the sites of planet formation. In the standard models of planet formation, the effect of the magnetic field is not seriously considered because the ionization degree is supposed to be very small at radii where the planet formation should take place. The low ionization is due to the high density of the midplane of the disk, where the recombination rate is too high to maintain ionization only through cosmic rays.

¹Department of Physics, Kyoto University, Kyoto 606-8502, Japan; inutsuka@tap.scphys.kyoto-u.ac.jp

²Institute of Laser Engineering, Osaka University, Suita, Osaka 565-0871, Japan; sano@ile.osaka-u.ac.jp

However, if the ionization degree is sufficiently high, the magnetorotational instability (MRI) is important and the disk becomes turbulent (Balbus & Hawley 1991, 1998; Stone et al. 2000). The combination of Maxwell and Reynolds stresses in magnetohydrodynamic turbulence provides an efficient mechanism for the radial transport of the angular momentum in the disk, which results in gas accretion onto the central star through the disk. Therefore, the determination of ionization degree in realistic protoplanetary disks is very important and remains the subject of extensive research (Sano et al. 2000; Glassgold et al. 2000; Fromang et al. 2002; Salmeron & Wardle 2003; Matsumura & Pudritz 2003). The existence of dust grains in the disk has a drastic effect on the ionization degree, because the recombination on the dust grains is very efficient in the high-density regime. Most of the studies on the ionization degree with the effect of dust grains show that there is a dead zone in which gas effectively decouples with the magnetic field. The location of the dead zone spans from a few tenths of AU to tens of AU from the central star depending on the (column) density distribution of the disk.

The existence of the dead zone should have important implications for the evolution of protoplanetary disks and the formation of planets (e.g., Gammie 1996, Matsumura & Pudritz 2005) and their migration in the disk (e.g., Terquem 2003, Laughlin, Steinacker, & Adams 2004, Nelson & Papaloizou 2004), which requires more detailed analysis on the validity of its existence and its precise location. In this Letter, we reanalyze the ionization degree taking into account additional sources of ionization that are not considered previously.

2. Ionization Required for MRI

Ionization Degree The linear stability analyses and nonlinear simulations show that the MHD turbulence in the disk is driven by the MRI if the magnetic Reynolds number is greater than about unity: $Re_M = v_A^2/(\Omega\eta) \gtrsim 1$ (Sano, Inutsuka, & Miyama 1998; Fleming, Stone, & Hawley 2000; Sano & Stone 2002a,b), where v_A is the Alfvén velocity and the resistivity η can be expressed as a function of the electron number density x_e and temperature T : $\eta \approx 2 \cdot 10^2 T^{1/2}/x_e$. Thus, the condition is expressed as

$$Re_M = \frac{v_A^2}{\Omega\eta} \approx \frac{C_s H}{\beta\eta} = \left(\frac{10}{\beta}\right) \left(\frac{x_e}{10^{-13}}\right) \left(\frac{H}{0.1\text{AU}}\right) \gtrsim 1, \quad (1)$$

where C_s denotes the sound speed, H is the scale height of the disk, and $\beta = 2C_s^2/v_A^2$ is the so called plasma β . In order to satisfy $Re_M > 1$, the number fraction of electrons should be larger than $x_e = 10^{-13}$ at $r \approx 1\text{AU}$ for typical protoplanetary disks with $\beta = 10$. Numerical simulations on the saturation level of the nonlinear development of MRI have shown that a typical value of β ranges from 10 to 100 (e.g., Hawley, Gammie, & Balbus 1995, Sano et al.

2004). In this Letter we adopt $\beta = 10$ as a fiducial value that corresponds to $B \approx 6$ Gauss for $n = 10^{15} \text{cm}^{-3}$ and $T \approx 3 \cdot 10^2$ K.

Scaling Relation in High Density Regime with Dust grains The fractional number densities of various charged species in the protoplanetary disks with dust grains are essentially determined by a single parameter that is the ionization rate ζ divided by the number density $n \approx 2n_{\text{H}_2}$ and calculated by Sano et al (2000). Figure 1 plots the fractional number densities of three important species as a function of ζ/n . The range of the fractional number density of electrons required for MRI, $x_e \approx 10^{-13}$, corresponds to the normalized ionization rate $\zeta/n \approx 3 \cdot 10^{-28} \text{sec}^{-1} \text{cm}^3$, i.e.,

$$\zeta \approx 3 \cdot 10^{-13} \left(\frac{n}{10^{15} \text{cm}^{-3}} \right) \text{sec}^{-1}. \quad (2)$$

This shows that for $n = 10^{15} \text{cm}^{-3}$ MRI requires an ionization rate 30,000 times higher than the standard cosmic-ray rate ($\zeta_{\text{CR}} \approx 10^{-17}$, see, e.g., Spitzer 1978). In the following, we consider whether this ionizing source is available or not.

3. Feedback from MHD turbulence

The column density of typical protoplanetary disks is so high that the external ionizing radiation (UV and X-rays) cannot penetrate deep into the midplane. We should consider the other mechanisms of ionization. Sano & Inutsuka (2001) have shown that the energy dissipation rate $Q [\text{erg cm}^{-3} \text{sec}^{-1}]$ in the turbulence driven by MRI in the disk can be calculated by the time and spatial average of the $r\phi$ -component of the stress tensor:

$$Q = \frac{3\Omega}{2} \left\langle \left\langle \rho v_R \delta v_\phi - \frac{B_R B_\phi}{4\pi} \right\rangle \right\rangle \approx 0.03 \Omega \langle \langle B^2 \rangle \rangle, \quad (3)$$

where the double angle brackets denote time and spatial average (see also Table 4 of Sano et al. 2004). Most of this energy dissipation is supposed to be used for heating of the gas. In a quasi-steady state condition, the input thermal energy is eventually converted to radiation that escapes from the disk. Here we propose that some fraction of the energy dissipation should be used for ionization.

Energy Budget for Sustaining Ionization First we consider whether the energy input available in the MHD turbulence can provide sufficient energy for the ionization required for MRI. The ratio f_{ionize} of the energy required for the ionization to the energy available in the

MRI-driven turbulence can be calculated as

$$f_{\text{ionize}} = \frac{\epsilon_{\text{ionize}} \zeta n}{Q} = 0.03 \left(\frac{n}{10^{15} \text{cm}^{-3}} \right)^2 \left(\frac{2\pi \text{yr}^{-1}}{\Omega} \right) \left(\frac{6\text{G}}{B} \right)^2 \left(\frac{\epsilon_{\text{ionize}}}{13.6\text{eV}} \right). \quad (4)$$

This is sufficiently small for typical conditions in protoplanetary disks even at 1 AU. Therefore, if the energy dissipation in MHD turbulence can be used for ionization of the gas, the positive feedback loop of MRI can maintain MHD turbulence in the protoplanetary disks as schematically shown in Figure 2.

4. Available Ionization Processes

The next question is about the microscopic processes of ionization: What is the process responsible for ionization in weakly ionized dusty gas?

Thermal Ionization Many numerical studies have shown that the saturated state of MRI-driven turbulence is the so-called high- β plasma: $\beta = 2C_s^2/v_A^2 \gtrsim 10$; i.e., the magnetic energy is smaller than the thermal energy of the gas. Therefore, even a rapid conversion of all the magnetic energy into thermal energy (via, e.g., magnetic reconnection) can raise the temperature of the gas only slightly. Thus, the magnetic dissipation does not seem to result in thermal ionization.

Note, however, that this argument requires great care because the thermal ionization of some alkali metals takes place at quite low temperature, $T \approx 1000\text{K}$ (e.g., Umebayashi 1983). We also note that the realistic saturation level of the magnetic field strength is still unknown and remains to be determined theoretically. Obviously, this requires more detailed understanding of the saturation mechanism of the MRI-driven turbulence, which includes the understanding of the physics of rapid magnetic reconnection in the case of a high- β environment (e.g., Sano et al. 2004).

Electron Mobility Let us assume that the thermal ionization is not promising, and consider another mechanism. In weakly ionized plasma, electrons have a considerable bulk velocity to maintain current and, hence, magnetic field. For sufficiently high density, the collision frequency of electrons is larger than the gyration frequency so that the conductivity tensor is isotropic and diagonal. Thus, we can assume that the average velocity of the electrons is essentially in the direction of the electric field ($-\vec{v}_e || \vec{E}$) in the comoving frame of the fluid. The averaged velocities of charged particles are related to the current density $\vec{j}$ via $\vec{j} = \sum_i e q_i n_i \langle \vec{v}_i \rangle$ or $e q_i n_i \langle v_i \rangle \equiv f_i j$, where the angle brackets mean a phase-space average,

and f_i is the fraction of the current due to the i th species of charged particle. From one of Maxwell's equations, $4\pi\vec{j} = c\nabla \times \vec{B}$, we estimate $j \approx cB/(2\lambda)$, where λ is the typical length scale of the magnetic field. Thus, we can estimate the average velocities of the charged species:

$$|\langle v_e \rangle| \approx \frac{cB}{2\lambda en_e} = 42 \left(\frac{B}{6\text{G}} \right) \left(\frac{0.03\text{AU}}{\lambda} \right) \left(\frac{10^{15}\text{cm}^{-3}}{n_H} \right) \left(\frac{10^{-13}}{x_e} \right) \text{km/s}, \quad (5)$$

where we may assume $f_e \approx 1$ for $\zeta/n \gtrsim 3 \cdot 10^{-28} \text{sec}^{-1} \text{cm}^3$ and $\lambda \lesssim 0.3H$ accords with the result of numerical simulations in Sano et al. (2004). This average electron velocity $\langle v_e \rangle$ is surprisingly large. Next we show its implication for the distribution function of electrons.

Energetic Electrons To clarify the possible ionization process, we should go into the details of microphysics. The electron distribution function $f(\vec{p})$ in weakly ionized plasma was already studied in detail (Druyvesteyn & Penning 1940; Landau & Lifshitz 1993).

If we choose the z -direction anti-parallel to $\vec{E}$, the electron distribution function is given by the following form in spherical coordinates (p, θ, ϕ) for the phase space:

$$f \approx f_0(p) + f_1(p) \cos(\theta), \quad (6)$$

$$f_0(p) = A \exp \left(-\frac{3\epsilon^2}{k_B^2 T^2 \gamma^2} \right), \quad (7)$$

$$f_1(p) = \frac{6\epsilon}{k_B T \gamma} \sqrt{\frac{m_e}{M}} f_0, \quad (8)$$

where

$$\epsilon = \frac{p^2}{2m_e}, \quad (9)$$

$$\gamma \equiv \frac{eEl}{k_B T} \sqrt{\frac{M}{m_e}} \gg 1, \quad (10)$$

and m_e is the electron mass and $M \approx 3670m_e$ is the hydrogen molecule mass. The normalization factor A is proportional to the number density of electrons. From these, we can calculate the average energy and the average velocity in the direction of the electric field.

$$\langle \epsilon \rangle = 0.43eEl \sqrt{\frac{M}{m_e}}, \quad \langle v_z \rangle = 0.9 \sqrt{\frac{eEl}{m_e}} \left(\frac{m_e}{M} \right)^{\frac{1}{4}} \quad (11)$$

$$\Rightarrow \langle \epsilon \rangle = 0.53M \langle v_z \rangle^2. \quad (12)$$

At first glance, it is surprising that the mass of the collision partner M appears in equation (12) instead of the electron mass m_e . This corresponds to the very low mobility of electrons

in the collisional medium, which is due to $m_e \ll M$. If we insert the expression for $\langle v_e \rangle \approx 42 \text{ km/sec}$ in equation (5) into $\langle v_z \rangle$ in equation (12), we have $\langle \epsilon \rangle = 19 \text{ eV}$, which is larger than the ionization potential of the hydrogen molecule. Thus, most of the free electrons have sufficient energies for ionization for this choice of parameters. This means that a significant fraction of the collisions may result in ionization, corresponding to a too high ionization rate. In reality, inelastic collisions with ionization/excitation losses should modify the distribution function (eq.[6]) that is obtained by considering only elastic scattering. A higher ionization degree results in a lower average velocity of electrons (see eq.[5]), which in turn reduces the ionizing collision according to equation (12). Therefore we expect that, in reality, the ionization degree might be on the order of $x_e \approx 10^{-13}$.

5. Yet Another Mechanism to Maintain Ionization Degree in Dust-Free Regions

The analysis of the ionization degree in the previous section is focused on the mid-plane of the disk, where external ionizing radiation is not available owing to the high column density of the disk. However, the gas density of the disk is a decreasing function of height (distance from the mid-plane), and thus, the surface layer of the disk is supposed to be well ionized by the incident interstellar radiation field, or by the radiation from the other stars in the cluster environment (see, e.g., Adams & Myers 2001). In the fully developed turbulence driven by MRI in the disk, the fluid elements travel from place to place according to eddies. The inverse cascade of turbulent energy tends to result in the state in which most of the power is in the largest eddies (Sano & Inutsuka 2001; Sano et al. 2004). This implies that each fluid element travels from the surface layer of the disk to the mid-plane, and vice versa. We should take into account this motion in the analysis of the ionization degree. For simplicity, let us consider the region with a very small amount of dust grains or the case in which the dust grains have grown up to larger sizes so that the total surface area of grains has become negligible. If we can ignore the recombination on the dust grains, the recombination should happen in the gas phase. The rate equation for the fractional electron number ($x_e = n_e/n$) is written as $dx_e/dt = \zeta - \sum_j \beta_j x_e n_j = \zeta - \beta' n x_e^2$, where β_j represents the coefficient for the recombination of electrons with the j th species of positively charged particles and $\beta' \equiv \sum_j \beta_j (x_j/x_e)$ is the effective recombination coefficient. Let us consider the fluid element that travels from the surface to the midplane. The approximate time evolution of the ionization degree $x_e(t)$ can be obtained by solving the rate equation with $\zeta = 0$ and the approximately constant β' :

$$x_e(t)^{-1} = x_{e,0}^{-1} + 10^{11} \left(\frac{\beta'}{3 \cdot 10^{-12} \text{ sec}^{-1} \text{ cm}^3} \right) \left(\frac{n}{10^{15} \text{ cm}^{-3}} \right) \left(\frac{t}{1 \text{ yr}} \right), \quad (13)$$

where $x_{e,0}$ is the initial ionization degree. For the high-density regime without the effect of grains, the recombination with positively charged metals is the most important process (Sano et al. 2000; Fromang et al. 2002). That is, within an eddy turnover time (≈ 1 yr at 1 AU), the ionization degree keeps the level of $x_e \gg 10^{-13}$ even without ionization due to cosmic rays, if we adopt the standard value of $\beta' \approx 3 \cdot 10^{-12} \text{cm}^3 \text{sec}^{-1}$ for metals and $x_{e,0} \gg 10^{-13}$. In other words, the ionized region penetrates into the neutral region by the turbulent motions. In this way, the turbulent motions efficiently homogenize the ionization degree if the recombination rate is sufficiently low. This spreading out of the weakly ionized region should be an important mechanism for activating MRI in the region that has initially low ionization. Note that this mechanism is not so effective in the region with the standard interstellar dust grains, because the recombination on the grain surface is much faster than in the gas phase.

6. Conclusions

Once the ionization degree becomes sufficiently high (eq.[1]), MRI-driven turbulence is the most promising mechanism for gas accretion in protoplanetary disks. Conversely, once the MRI-driven turbulence becomes active, it can consistently maintain the ionization degree without help of other external ionizing sources: the macroscopic energy dissipation in turbulence provides sufficient energy for the ionization that is required for activating MRI, and microscopically the energetic electrons in weakly ionized gas should provide a sufficient rate of collisional ionization. The turbulent spreading of the ionized gas also provides the mechanism of spreading the magnetorotationally unstable region. Altogether we conclude that most of the region in the protoplanetary disk should be sufficiently ionized, thus, effectively removing the dead zone. The absence of the dead zone in the protoplanetary disks has tremendous impact on the theoretical modeling of planet formation.

The authors thank the anonymous referee for the prompt and useful comments. They also thank K. Shibata and T. Nakano for valuable discussions. The authors are supported by the Grant-in-Aid (15740118, 16077202, 16740111, 17039005) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan. This work is supported by the Grant-in-Aid for the 21st Century COE “Center for Diversity and Universality in Physics” from MEXT of Japan.

REFERENCES

- Adams, F. C. & Myers, P. C. 2001, ApJ 553, 744
- Balbus, S. A., & Hawley, J. F. 1991, ApJ, 376, 214
- Balbus, S. A., & Hawley, J. F. 1998, Rev. Mod. Phys. 70 1
- Druyvesteyn, M. J. & Penning, F. M. 1940, Reviews of Modern Physics 12, 87
- Fleming, T. P., Stone, J. M., & Hawley, J. F. 2000, ApJ, 530, 464
- Fromang, S., Terquem, C., & Balbus, S. 2002, MN 329, 18
- Gammie, C. F. 1996, ApJ, 457, 355
- Glassgold, A. E., Feigelson, E. D., & Montmerle, T. 2000, in Protostars and Planets IV, ed. V. Mannings, A. P. Boss, & S. S. Russell (Tucson: Univ. Arizona Press), 429
- Hawley, J. F., Gammie, C. F., & Balbus, S. A. 1995, ApJ, 440, 742
- Landau, L. D. & Lifshitz, E. M. 1993, Physical Kinetics (New York: Pergamon)
- Laughlin, G., Steinacker, A., & Adams, F. C. 2004, ApJ 608, 489
- Matsumura, S. & Pudritz, R. 2003, ApJ 598, 645
- Matsumura, S. & Pudritz, R. 2005, ApJ 618, L137
- Nelson, R. P. & Papaloizou, J. C. B. 2004, MNRAS 350, 849
- Salmeron, R. & Wardle, M. 2003, MN 345, 992
- Sano, T., Inutsuka, S., & Miyama, S. M. 1998, ApJ 506, L57
- Sano, T. & Inutsuka, S. 2001, ApJ 561, L179
- Sano, T., Miyama, S. M., Umebayashi, T., & Nakano, T. 2000 ApJ 543, 486
- Sano, T. & Stone, J. M. 2002, ApJ 570, 514
- Sano, T. & Stone, J. M. 2002, ApJ 577, 534
- Sano, T., Inutsuka, S., Turner, N. J., & Stone, J. M. 2004, ApJ 605, 321
- Spitzer, L. Jr. 1978, Physical Processes in the Interstellar Medium (New York: Wiley)

Stone, J. M., Gammie, C. F., Balbus, S. A., & Hawley, J. F. 2000, in *Protostars and Planets IV*, ed. V. Mannings, A. P. Boss, & S. S. Russell (Tucson: Univ. Arizona Press), 589

Terquem, C. E. J. M. L. J. 2003 *MNRAS* 341, 1157

Umebayashi, T. 1983, *Prog. Theor. Phys.* 68, 480




Fig. 1.— Fractional number densities of electrons (e^-), metals (M^+), and molecules (mo^+) as a function of the ionization rate divided by the total number density in the model with dust grains. Recombinations on grain surfaces are taken into account as well as the radiative and dissociative recombinations in the gas phase. The size of the dust grain is assumed to be $0.1 \mu\text{m}$. See Sano et al. (2000) for details.




Fig. 2.— Schematic diagram for the feedback loop in the turbulence driven by magnetorotational instability (MRI). Once the protoplanetary disk is sufficiently ionized, the MRI occurs and the magnetohydrodynamic turbulence develops and saturates. At the saturated state, most of the energy dissipation results in the heating of the gas and its conversion into the radiation that eventually escapes from the disk. If only a small fraction of the energy dissipation is used for ionization, the disk keeps this magnetic activity (see eq.[4]). One of the possible processes of ionization should be the collision between the energetic electrons and neutral particles (see Section 4).